

Contains a source of live (viable) naturally occurring microorganisms.

GUARANTEED ANALYSIS: (per pound)

Crude Protein (minimum)	2 %
Crude Fat (minimum)	1 %
Crude Fiber (maximum)	11 %
Sodium (minimum)	4.87 %
Sodium (maximum)	5.87 %
Potassium (minimum)	1.38 %

Live Microbial Count (minimum)

20 Billion Colony Forming Units

(Lactobacillus acidophilus, Lactobacillus casei, Enterococcus faecium, Lactobacillus plantarum, Lactobacillus lactis, Pediococcus acidilactici, Bifidobacterium bifidum, Bifidobacterium infantis, Bifidobacterium longum, Bacillus licheniformis, Bacillus subtilis)

INGREDIENTS:

Roughage products, reed-sedge peat, sodium bicarbonate, sodium chloride, hydrolyzed yeast, guar gum, dextrose, inulin, dried kelp, potassium chloride, vitamin A supplement, vitamin D3 supplement, vitamin E supplement, thiamine mononitrate, riboflavin, niacin, calcium pantothenate, pyridoxine hydrochloride, folic acid, biotin, vitamin B12 supplement, ascorbic acid, sodium silico aluminate, Yucca schidigera plant extract, dried Lactobacillus acidophilus fermentation product, dried Lactobacillus casei fermentation product, dried Enterococcus faecium fermentation product, dried Lactobacillus plantarum fermentation product, dried Lactobacillus lactis fermentation product, dried Pediococcus acidilactici fermentation product, dried Bifidobacterium bifidum fermentation product, dried Bifidobacterium infantis fermentation product, dried Bifidobacterium longum fermentation product, dried Bacillus subtilis fermentation product, dried Bacillus licheniformis fermentation product.

RECOMMENDED USES:

Designed as a nutritional supplement for calves at shipping, during fluid loss and when diarrhea is present. Supplies dietary fibers, electrolytes and beneficial microbial cultures.

DIRECTIONS:

(1 scoop = 1 ounce) For calves with fluid deficiency, continue feeding milk at alternate times (at least 30 minutes) after feeding PectiLyte with water.

Consult veterinarian for proper drenching/tubing procedures.

DAILY MAINTENANCE: Mix 1 oz. PectiLyte / 2 quarts warm water, milk or milk replacer. Feed once or twice daily.

AT RECEIVING: Mix 1 to 2 oz. PectiLyte / 2 quarts warm water once or twice daily at alternate times to milk or milk replacer.

SLIGHT FLUID DEFICIENCY: Mix 2 to 4 oz. PectiLyte / 2 quarts warm water twice daily.

MODERATE FLUID DEFICIENCY: Mix 4 oz. PectiLyte / 2 quarts warm water two to three times daily.

STORAGE: Keep sealed between uses. Store in cool, dry place.

PRODUCT CODE

GN-01120	11 pound pouch (4 pouches / case)
GN-00125	50 pound bags (1 / case)
GN-00125	1 ton (50 pound bags, 40 / pallet)



TS_PectiLyteRevised02052021

Bio-Vet, Inc. • 300 Ernie Drive • Barneveld, WI 53507, U.S.A. • (800) 246- 8381 • Fax (608) 924- 7006 • www.Bio-Vet.com